



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## MECHANICS.

When this issue was made up, solutions had been received for 285-8-9 and 292. Solutions of 266, 268, 269, 271 to 275, inclusive, and 277 to 279 inclusive are desired.

**293. Proposed by B. F. FINKEL, Drury College.**

A man of weight  $W$  stands on smooth ice; prove that if, when he gradually parts his legs, kept straight, with his feet in contact with the ice, the pressure of his feet on the ice be constant, his head will descend with uniform acceleration; and that, if  $f$  be the acceleration of his head, when his feet exert no pressure on the ice, their pressure on the ice, were  $f'$  the acceleration of his head, would be equal to  $\frac{f-f'}{f} W$ .

(From WALTON'S *Problems in Theoretical Mechanics*.)

**294. Proposed by EMMA GIBSON, Drury College.**

A sphere, revolving about a diameter and not acted on by any extraneous force, expands symmetrically; prove that its vis viva varies inversely as its moment of inertia about a diameter.

## NUMBER THEORY.

When this issue was made up, solutions had been received for 206-7, 210, 213. Solutions of 192 and 196 are desired.

**215. Proposed by R. D. CARMICHAEL, Indiana University.**

Find one or more values of  $n$  such that a polygon of  $n$  sides shall have the number of its diagonals equal to the cube of an integer.

**216. Proposed by ELIJAH SWIFT, Princeton, N. J.**

If  $p$  is a prime  $> 3$ , show that  $\sum_{a=1}^{a=p-1} 1/a \equiv 0 \pmod{p^2}$ , where  $1/a \equiv x$ , if  $ax \equiv 1 \pmod{p^2}$

**217. Proposed by E. T. BELL, Seattle, Wash.**

- (i) If  $r$  is a prime greater than 2, and  $p \equiv 2^r + 1$  is prime, the only solution, when  $n$  is greater than 2, of  $x^n - y^n = p$ , is  $n = 3, x = 2, y = 1$ .
- (ii) The only primes that are simultaneously of the forms  $4k + 1$  and  $3^m - 2^m$  are 1 and 5.
- (iii) Generalize (ii).

## SOLUTIONS OF PROBLEMS.

## ALGEBRA.

**385. Proposed by J. F. LAWRENCE, Stillwater, Oklahoma.**

Show that if  $p$  is prime and  $> 3$ ,  $(2p)! - 2 \cdot p!p!$  is divisible by  $p^5$ .

SOLUTION BY ELIJAH SWIFT, Princeton, N. J.

Dividing the above expression by  $2p \cdot p!$  we have to show that

$$(1) \quad (2p-1)(2p-2)(2p-3) \cdots (2p-(p-1)) - (p-1)!$$

is divisible by  $p^3$ .

Consider the polynomial  $P(x) = (x-1)(x-2)(x-3) \cdots (x-p+1)$ .<sup>1</sup> If this is expanded in powers of  $x$ , we obtain an expression of the form

$$x^{p-1} - A_1 x^{p-2} + A_2 x^{p-3} + \cdots + A_{p-3} x^2 - A_{p-2} x + A_{p-1},$$

<sup>1</sup> See BACHMANN, *Niedere Zahlentheorie*, p. 155. I have changed signs, but evidently the same results hold.